

Mira B Rejuvenated¹

Brian E. Wood² and Margarita Karovska³

ABSTRACT

Recent ultraviolet spectra from the *Hubble Space Telescope* (HST) and *Far Ultraviolet Spectroscopic Explorer* (FUSE) satellites demonstrate that UV line and continuum fluxes observed from Mira B are increasing back towards the levels that the *International Ultraviolet Explorer* observed in 1979–1980 and 1990–1995, after having been found in a very low state by HST and FUSE in 1999–2001. The UV emission is associated with accretion of material onto Mira B from Mira A’s massive wind, so the variability is presumably due to variations in accretion rate. From wind absorption features, we estimate a Mira B mass loss rate of $2.5 \times 10^{-12} M_{\odot} \text{ yr}^{-1}$, indicating that Mira B’s wind has increased in strength along with the accretion rate. The UV variability may be associated with a previously reported 14-year periodicity in Mira B’s optical emission.

Subject headings: accretion, accretion disks — binaries: close — stars: individual (o Ceti) — stars: winds, outflows — ultraviolet: stars

1. INTRODUCTION

Mira AB is the nearest symbiotic system at a distance of only ~ 130 pc, (Perryman et al. 1997), and the only symbiotic binary in which the components of the system have been resolved at wavelengths ranging from X-rays to radio (Karovska et al. 1997, 2005; Matthews & Karovska 2006). The primary (Mira, o Ceti, HD 14386) is a very large pulsating AGB star (photospheric diameter of $\sim 500 R_{\odot}$), with a massive, cool wind of $\sim 10^{-7} M_{\odot} \text{ yr}^{-1}$

¹Based on observations with the NASA/ESA Hubble Space Telescope and the NASA-CNES-CSA Far Ultraviolet Spectroscopic Explorer. The former observations were obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. FUSE is operated for NASA by the Johns Hopkins University under NASA contract NAS5-32985.

²JILA, University of Colorado, Boulder, CO 80309-0440; woodb@origins.colorado.edu.

³Smithsonian Astrophysical Observatory, 60 Garden St., Cambridge, MA 02138; mkarovska@cfa.harvard.edu.

(Bowers & Knapp 1988). The companion (Mira B, VZ Ceti) is accreting material from Mira A, but the exact nature of the accretor is uncertain (Jura & Helfand 1984); it has usually been assumed to be a white dwarf (Joy 1954; Karovska et al. 2005). The current angular separation between the components is $\sim 0.6''$ (Karovska et al. 1997). The orbit of the binary is poorly known, but Prieur et al. (2002) estimate a long orbital period of ~ 500 years, and this orbit suggests that the distance from Mira A to Mira B is close to the apparent separation in the plane of the sky, about 70 AU. This interacting binary, being one of the few accretion systems in which the components of the system are resolvable, provides a unique opportunity to study the spectral energy distributions of the components of an interacting binary individually, as well as being a laboratory for testing accretion models.

In 1995, the Faint Object Camera (FOC) instrument on the *Hubble Space Telescope* (HST) resolved the Mira system for the first time spatially and spectrally at UV and optical wavelengths (Karovska et al. 1997). Since then, the system has been monitored from UV to radio wavelengths. In the past few years, we have been witnessing dramatic changes in Mira B, especially at UV wavelengths (Karovska et al. 1997; Wood et al. 2001). Spectra from the Space Telescope Imaging Spectrograph (STIS) instrument on HST imply a substantial decrease in accretion luminosity in the late 1990s, compared with fluxes observed at earlier times by the *International Ultraviolet Explorer* (IUE) and HST/FOC. Furthermore, a forest of Ly α -fluoresced H₂ emission lines dominated the HST spectra of Mira B in 1999, despite not being seen at all in any previous IUE or HST spectra (Wood et al. 2001, 2002). Observations from the *Far Ultraviolet Spectroscopic Explorer* (FUSE) demonstrated that the low accretion luminosity and prominent H₂ emission persisted into 2001 (Wood & Karovska 2004). On 2003 December 6, *Chandra* observations detected an unprecedented soft X-ray outburst from Mira A (Karovska et al. 2005), further evidence of remarkable dynamicism in the Mira AB system. The outburst was also detected in contemporaneous H α observations and is likely associated with a magnetic flare followed by a mass ejection or jet-like activity (Karovska et al. 2005, Karovska 2006, in prep.).

We here present recent UV observations of Mira B from HST and FUSE, obtained following the X-ray outburst. We compare these data with past observations by these satellites and by IUE. We discuss the long term variations of Mira B and a possible correlation with long term optical variability.

2. UV SPECTROSCOPY OF MIRA B FROM 1979–2001

Numerous UV spectra of Mira AB were obtained in 1979–1980 and 1990–1995 by IUE. This data set includes 32 spectra taken in IUE’s short-wavelength, low-resolution mode (i.e.,

SW-LO spectra, covering $\sim 1150 - 1950 \text{ \AA}$). These far-UV (FUV) spectra are dominated by broad, hot lines such as C II $\lambda 1335$, Si IV $\lambda 1400$, C IV $\lambda 1550$, Si III] $\lambda 1892$, and C III] $\lambda 1908$, which are produced by material accreting onto Mira B from Mira A’s massive wind. The few high-resolution spectra of this wavelength region (i.e., SW-HI spectra) are very noisy, but they suggest very broad widths for these lines, consistent with emission from a rapidly rotating accretion disk (Reimers & Cassatella 1985). Unlike IUE, HST can resolve the binary, and UV spectra from HST confirm that the broad lines are from Mira B, and that Mira A contributes very little to the IUE spectra below $2000 \text{ \AA}$ (Karovska et al. 1997; Wood et al. 2001).

Only a modest level of UV variability is present within the large IUE SW-LO data set, with line and continuum fluxes varying by about a factor of 2. However, when HST observed Mira B in 1999 August using STIS, the character of Mira B’s FUV spectrum had changed dramatically. The UV continuum and all of the aforementioned UV emission lines had weakened by at least a factor of 20, and the FUV spectrum was now dominated by many narrow H_2 lines fluoresced by H I $\text{Ly}\alpha$ (Wood et al. 2001), which are not detectable at all in the IUE spectra. The accretion onto Mira B results in a warm, fast wind from the star that is most clearly apparent in broad absorption features in the Mg II h & k lines at $2800 \text{ \AA}$. Given the presence of accretion, this wind is envisioned to be of a bipolar nature. Analysis of the Mg II absorption demonstrates that the substantial drop in UV flux from Mira B seen by HST is accompanied by a corresponding drop in its mass loss rate (Wood et al. 2002).

The low UV fluxes observed by HST in 1999 are most naturally explained by a substantial decrease in the accretion rate onto Mira B, which also led to a weaker wind from the star. The sudden prominence of the H_2 lines in the HST spectra suggests that unlike other UV lines, the H I $\text{Ly}\alpha$ line responsible for fluorescing these lines had *not* decreased in strength relative to the IUE era. Studying $\text{Ly}\alpha$ in IUE spectra is very difficult due to geocoronal contamination, but Wood et al. (2002) still uncovered evidence to support the contention that $\text{Ly}\alpha$ fluxes are similar in the HST and IUE spectra.

A 2001 November observation from FUSE found Mira B to be still in a very low state, with C III $\lambda 1175$ fluxes comparable to those in the 1999 HST/STIS spectrum (Wood & Karovska 2004). Like the STIS data, this FUSE spectrum also shows numerous narrow H_2 lines. Ground-based observations have suggested that Mira B’s optical light curve may have a period of 14 years, with the last published minimum being in 1971 (Joy 1954; Yamashita & Maehara 1977). Wood & Karovska (2004) noted that the numerous IUE SW-LO observations are all near predicted maxima of this cycle, while the 1999 HST and 2001 FUSE observations are near a predicted minimum. Thus, it is possible that the observed long-term UV variability could be correlated with previously detected optical variations. If

the optical periodicity is real and if a connection between the optical and UV light curves truly exists, Mira B’s UV fluxes should increase towards a 2006–2007 maximum predicted by the 14-year optical periodicity. The new 2004 HST and FUSE spectra presented here allow us to test this prediction, and to see if the character of Mira B’s UV spectrum has changed once again.

3. RECENT HST AND FUSE OBSERVATIONS

3.1. Spectral Line Variability

Mira B was observed by HST with the STIS instrument on 2004 February 16. A near-UV (NUV) spectrum from 2274–3119 Å was taken using the moderate resolution E230M grating, with an exposure time of 600 s, and Mira B’s FUV spectrum from 1140–1729 Å was observed using the E140M grating, with an exposure time of 2100 s. Both spectra were taken through the $0.2'' \times 0.2''$ aperture.

Mira was also observed by FUSE on 2004 August 31. This was an 8603 s exposure through the LWRS aperture, covering the 905–1187 Å spectral range. This represents only a fraction of the requested 30 ksec exposure, so the new FUSE spectrum is much noisier than the 2001 FUSE observation (Wood & Karovska 2004). A C III $\lambda 1175$ flux can still be measured from the data and some of the strongest H₂ lines can still be discerned.

Figures 1 and 2 compare UV spectral regions observed by HST in 2004 with the old 1999 spectra. Five of the Figure 1 panels focus on spectral regions containing FUV emission lines of H I, O I, C II, Si IV, and C IV. Some of these lines are blended with H₂ emission, so the locations of strong, narrow H₂ lines are also indicated in the figure. The last panel shows the NUV spectrum in the region of the Fe II UV1 multiplet lines. The line locations indicated in Figure 1 are with respect to Mira’s rest frame, where the rest wavelengths are taken from Abgrall et al. (1993) and Morton (2003), and we assume a radial velocity of 56 km s^{−1} for the Mira system (Bowers & Knapp 1988; Planesas et al. 1990; Josselin et al. 2000).

Figures 1 and 2 show that UV fluxes from Mira B have increased dramatically since 1999, with the O I, C II, Si IV, C IV, and Mg II lines being much stronger than they were. The notable exceptions to this rule, the H I Ly α line and the H₂ lines fluoresced by Ly α , will be discussed further in §3.3. The UV continuum is also much stronger in 2004 (see the Fe II panel in Fig. 1). In Figure 3, we plot measured fluxes as a function of time for many different lines, including measurements from IUE, HST, and FUSE. For C III $\lambda 1175$, O I $\lambda 1300$, C II $\lambda 1335$, Si IV $\lambda 1400$, and C IV $\lambda 1550$, the IUE fluxes are measured from the 32

IUE SW-LO observations mentioned in §2. Examples of these spectra have been previously presented by Reimers & Cassatella (1985) and Wood et al. (2001). Note that the fluxes in Figure 3 are the total flux of these multiplets (e.g., for Si IV the flux includes both the $\lambda 1393$ and $\lambda 1402$ lines), given that the individual components of the multiplets are not resolved in the IUE SW-LO spectra.

Blended H_2 lines are removed from the data before measuring fluxes from the HST and FUSE data. This cannot be done for the low resolution IUE spectra, but thanks to the higher line fluxes seen by IUE, H_2 is not as strong a contaminant in the IUE data, so the correction is not as important (Wood et al. 2001). The C III $\lambda 1175$ line is the only line observable by all three satellites, though in the HST/STIS spectra and the 2004 FUSE spectrum the C III line is observed at very low S/N. In the higher quality 2001 FUSE spectrum, the H_2 lines add 1.6×10^{-14} ergs cm $^{-2}$ s $^{-1}$ to the C III feature (Wood & Karovska 2004), so we simply subtract this flux from each C III flux measurement to correct for the strong H_2 contamination in C III.

The IUE’s inability to resolve the Mira binary is generally not a problem since Mira A does not contribute significantly to most UV emission lines. However, Mira A can contribute to Mira AB’s Mg II lines at certain pulsation phases (Wood & Karovska 2000). Fortunately, in long-wavelength, high-resolution (LW-HI) IUE spectra, Mira A’s narrow Mg II lines are mostly contained within the saturated core of Mira B’s Mg II wind absorption. Therefore, the Mira A Mg II line can be removed before measuring a Mg II flux for Mira B. The IUE Mg II fluxes plotted in Figure 3 are all from LW-HI spectra. We do not use LW-LO spectra because of potential Mira A contamination. The Mg II fluxes in Figure 3 include both the h and k lines.

Figure 3 shows the dramatic decrease in UV line fluxes observed in 1999–2001. The highest temperature line, C IV $\lambda 1550$, exhibits a particularly large decrease of nearly a factor of 100. The fluxes increase significantly in 2004, but they are still lower than fluxes observed by IUE in 1979–1980 and 1990–1995. As discussed in §2, a 14-year periodicity has been detected in Mira B’s optical light curve (Joy 1954; Yamashita & Maehara 1977), and dotted lines in Figure 3 indicate the phase and period of this light curve. The increase in UV flux that we see in our 2004 spectra is roughly consistent with the predictions of the optical light curve, assuming that the UV emission follows the same periodic behavior as that inferred for the optical light. It therefore remains a plausible hypothesis that Mira B’s long-term UV variability is associated with the previously detected periodic optical variability. However, truly verifying this connection would require UV observations over at least another cycle.

3.2. Wind Absorption

The blue sides of all the resonance lines in Figures 1 and 2 are suppressed by absorption from Mira B’s wind, with the exception of C IV (see below). The wind absorption is more readily apparent in the new 2004 data than in the old 1999 spectra. This is in part due to much better S/N in the new data thanks to the higher fluxes, but it is also due to the wind absorption being intrinsically stronger, which is particularly evident for the Fe II and Mg II lines. The mass loss rate of Mira B’s wind can be estimated most easily from the Mg II wind absorption profiles, which have the highest S/N.

From analyses of the Mg II k line at 2796 Å, we have previously estimated a mass loss rate of $\dot{M} = 5 \times 10^{-13} \text{ M}_{\odot} \text{ yr}^{-1}$ from the 1999 HST spectrum, and $\dot{M} = 1 \times 10^{-11} \text{ M}_{\odot} \text{ yr}^{-1}$ from a typical IUE spectrum (Wood et al. 2002). Using the exact same methodology, we have fitted a wind absorption profile to the 2004 Mg II k line in Figure 2. The observed absorption profile is not smooth, suggesting a more complicated and structured wind than we can fit with such a simple model. Nevertheless, the mass loss rate inferred by the fit should still be a reasonable estimate. This fit suggests a mass loss rate of $\dot{M} = 2.5 \times 10^{-12} \text{ M}_{\odot} \text{ yr}^{-1}$ and a wind terminal velocity of $V_{\infty} = 450 \text{ km s}^{-1}$. For comparison, the absorption predicted by the wind absorption profile from 1999 is also shown in Figure 2, assuming $\dot{M} = 5 \times 10^{-13} \text{ M}_{\odot} \text{ yr}^{-1}$ and $V_{\infty} = 250 \text{ km s}^{-1}$. The wind absorption is significantly stronger in 2004, meaning that Mira B’s wind has clearly strengthened, but the mass loss rate is still not as high as the $\dot{M} = 1 \times 10^{-11} \text{ M}_{\odot} \text{ yr}^{-1}$ value estimated for the IUE era. The mass loss rate variations are consistent with the accretion rate variations inferred from the UV fluxes (see Fig. 3). Clearly, Mira B’s accretion rate is highly correlated with the strength of its wind.

The only atomic lines in Figure 1 that do not definitively possess wind absorption are the C IV lines. Small decreases in flux blueward of the center of the two C IV lines may indicate the presence of *some* C IV wind absorption, but this absorption is not nearly as prominent as the saturated absorption seen for the other lines. The presence of strong wind absorption for the Si IV $\lambda 1392$ line, and the relative lack of absorption for C IV implies a maximum temperature for the wind between $T = 6.0 \times 10^4 \text{ K}$ (Si IV’s line formation temperature) and $T = 1.0 \times 10^5 \text{ K}$ (C IV’s line formation temperature).

3.3. The Anomalous Ly α Line

The one clearly anomalous line in Figure 1 is H I Ly α . Although the 2004 Ly α flux is slightly higher than in 1999, this line does not show the dramatic flux increase that the

other emission lines show. The singular lack of variability for this line was also noted by Wood et al. (2002), who inferred that the $\text{Ly}\alpha$ flux during the IUE era cannot have been much stronger than in 1999, unlike all the other lines. The lack of variability for $\text{Ly}\alpha$ means that the H_2 lines that are fluoresced by $\text{Ly}\alpha$ are also relatively constant, as shown by the H_2 lines blended with C II and Si IV in Figure 1.

It is difficult to understand why the $\text{Ly}\alpha$ line behaves so differently from the continuum and all other emission lines, including lines formed at similar temperatures (O I, C II, and Mg II). Does this mean that the $\text{Ly}\alpha$ emission is coming from a different place than the other lines? If so, why? Another possibility is that the $\text{Ly}\alpha$ flux actually *does* vary along with the other lines, but the extremely high opacity of Mira B’s wind to $\text{Ly}\alpha$ moderates the variation. Wind absorption explains why the observed $\text{Ly}\alpha$ emission is entirely redshifted relative to line center (see Fig. 1). Analogous absorption is also seen for many of the other lines, but wind opacity will be orders of magnitude higher for $\text{Ly}\alpha$ since hydrogen is much more abundant than other atomic species.

Given that increases in UV flux are clearly accompanied by increases in wind strength, perhaps the enhanced wind obscuration largely cancels the flux increase for $\text{Ly}\alpha$. This hypothesis would require that $\text{Ly}\alpha$ photons are actually destroyed rather than just scattered by the wind. Perhaps this destruction could be accomplished by scattering into Mira B’s accretion disk, or extinction by dust entrained within Mira B’s wind.

4. CAUSES OF LONG-TERM VARIABILITY

The fundamental cause of Mira B’s substantial variability remains uncertain. Variable mass loss from Mira A could have caused the changes in the UV flux of Mira B observed in the past 10 years, signaling that the system is undergoing important transformations. These transformations may have set the stage for the remarkable X-ray outburst from Mira A that was observed by *Chandra* on 2003 December 6 (Karovska et al. 2005).

If the variability is associated with the possible 14-year optical periodicity, the reason for the periodicity is a mystery. The UV lines in particular are believed to originate from accretion processes on Mira B, so a periodicity in these lines would suggest a periodicity for the accretion rate. Accretion instabilities could easily lead to variability, but it is not clear why they would be periodic. Perhaps the gravitational influence of an undetected stellar or planetary companion could lead to a corresponding periodicity in the accretion rate, perhaps through wind focusing effects. Hints of a third body in the Mira AB system have been previously reported in the literature (Baize 1980; Karovska 1992; Karovska et al. 1993).

Since Mira A’s wind is the source of the accreting material, a periodicity in Mira A’s mass loss rate could in principle also lead to periodicity in the accretion rate. No such periodicity has yet been found in the light curve of Mira compiled by AAVSO, but searches for very low amplitude, long-term periodicity are ongoing (M. Templeton, private communication).

We hope to explore these possibilities further in the future, as we reach the expected maximum in 2006–2007 and over the next 14-year cycle, using different observational techniques including optical/IR interferometry. However, future UV high resolution spectra are very much in doubt due to the failure of the STIS instrument in 2004 August, and the reaction wheel and gyro failures on FUSE that currently limit observations to high declinations. Thus, new UV instrumentation may be necessary to continue spectroscopic observations in the ultraviolet.

This work was supported by NASA grant NAS8-39073 to the Smithsonian Astrophysical Observatory and NASA grant NNG05GC14G to the University of Colorado. MK is a member of the Chandra X-ray Center, which is operated by the Smithsonian Astrophysical Observatory under contract to NASA NAS8-39073.

REFERENCES

- Abgrall, H. A., Roueff, E., Launay, F., Roncin, J. -Y., & Subtil, J. -L. 1993, *A&AS*, 101, 273
- Baize, P. 1980, *A&AS*, 39, 86
- Bowers, P. F., & Knapp, G. R. 1988, *ApJ*, 332, 299
- Josselin, E., Maun, N., Planesas, P., & Bachiller, R. 2000, *A&A*, 362, 255
- Joy, A. H. 1954, *ApJS*, 1, 39
- Jura, M., & Helfand, D. J. 1984, *ApJ*, 287, 785
- Karovska, M. 1992, in *Complementary Approaches to Double and Multiple Star Research*, ed. H. A. McAlister & W. I. Hartkopf (ASP Conf. Ser., Vol. 32), 558
- Karovska, M., Hack, W., Raymond, J., & Guinan, E. 1997, *ApJ*, 482, L175
- Karovska, M., Nisenson, P. & Beletic, J. 1993, *ApJ*, 402, 311
- Karovska, M., Schlegel, E., Hack, W., Raymond, J. C., & Wood, B. E. 2005, *ApJ*, 623, L137
- Matthews, L. D., & Karovska, M. 2006, *ApJ*, 637, L49
- Morton, D. C. 2003, *ApJS*, 149, 205
- Perryman, M. A. C., et al. 1997, *A&A*, 323, L49
- Planesas, P., Bachiller, R., Martin-Pintado, J., & Bujarrabal, V. 1990, *ApJ*, 351, 263
- Prieur, J. L., Aristidi, E., Lopez, B., Scardia, M., Mignard, F., & Carbillet, M. 2002, *ApJS*, 139, 249
- Reimers, D., & Cassatella, A. 1985, *ApJ*, 297, 275
- Wood, B. E., & Karovska, M. 2000, *ApJ*, 535, 304
- Wood, B. E., & Karovska, M. 2004, *ApJ*, 601, 502
- Wood, B. E., Karovska, M., & Hack, W. 2001, *ApJ*, 556, L51
- Wood, B. E., Karovska, M., & Raymond, J. C. 2002, *ApJ*, 575, 1057
- Yamashita, Y., & Maehara, H. 1977, *PASJ*, 29, 319




Fig. 1.— A comparison of UV spectra of Mira B taken in 1999 (dotted line) and 2004 (solid line). The spectra are smoothed slightly for the sake of appearance. Dashed lines indicate the rest wavelengths (in Mira’s rest frame) of the line or lines indicated by the panel labels (e.g., H I Ly α , O I, C II, Si IV, C IV, and Fe II). Dot-dashed lines indicate the locations of strong, narrow H₂ lines.




Fig. 2.— A comparison of Mira B’s Mg II k line profile from 1999 (dotted histogram) and 2004 (solid histogram), plotted on a velocity scale in Mira B’s rest frame. A wind absorption profile has been fitted to the 2004 data (thick solid line), which suggests a mass loss rate of $\dot{M} = 2.5 \times 10^{-12} M_{\odot} \text{ yr}^{-1}$ and a terminal velocity of $V_{\infty} = 450 \text{ km s}^{-1}$. Also shown is the absorption predicted by a fit to the 1999 wind absorption (dot-dashed line), assuming $\dot{M} = 5 \times 10^{-13} M_{\odot} \text{ yr}^{-1}$ and $V_{\infty} = 250 \text{ km s}^{-1}$ (Wood & Karovska 2004). This greatly underpredicts the amount of absorption observed in 2004, illustrating the increase in wind strength from 1999 to 2004.




Fig. 3.— Fluxes of various UV emission lines as a function of time, based on spectra from IUE (triangles), HST (boxes), and FUSE (diamonds). The fluxes are normalized to the average value observed by IUE. The lines and normalization factors (in units of $\text{ergs cm}^{-2} \text{s}^{-1}$) are C III $\lambda 1175$: 4.1×10^{-13} , O I $\lambda 1300$: 8.1×10^{-13} , C II $\lambda 1335$: 6.5×10^{-13} , Si IV $\lambda 1400$: 5.5×10^{-13} , C IV $\lambda 1550$: 1.7×10^{-12} , and Mg II $\lambda 2800$: 1.3×10^{-11} . The dotted lines are schematic representations of the 14 year periodicity of Mira B suggested by optical observations (Joy 1954; Yamashita & Maehara 1977).